

PROCEEDINGS OF THE SIXTH
INTERNATIONAL TOBACCO SCIENTIFIC
CONGRESS

ACTES DU SIXIEME
CONGRES SCIENTIFIQUE INTERNATIONAL
DU TABAC

TOKYO
November, 14 to 20
1976

Organized by CORESTA and
THE JAPAN TOBACCO & SALT PUBLIC CORPORATION

PM3006448059

CONTENTS

TABLE DES MATIERES

First Part – The Congress Days Première Partie – Les Journées du Congrès

	Page
President of Congress – Président du Congrès	17
Organizing Committee – Comité d'Organisation	18
Council – Conseil	19
Scientific Commission – Commission Scientifique	20
Reading Committee – Comité de Lecture	21
Chairmen of Scientific Meetings – Présidents de Réunions Scientifiques	22
Opening Session – Session d'Ouverture	23
Closing Session – Session de Clôture	29

Second Part – Congress Proceedings Seconde Partie – Les Travaux du Congrès

Invited Papers – Communications Sollicitées

Kenneth R. KELLER Tobacco – the plant <i>Tabac – la plante</i>	39
Karl H. WEBER Recent changes in tobacco products and their acceptance by the consumer <i>Changements récents dans les produits du tabac et leur acceptation par le consommateur</i>	47
Sadayuki F. TAKAGI Ten chapters on olfaction <i>Dix chapitres sur l'olfaction</i>	64

Joint Meeting – Réunion Commune

SMITH H. H.	
Genetic engineering with tobacco protoplasts	75
<i>Engineering génétique sur des protoplastes du tabac</i>	
TSO T. C., GORI G. B.	
A novel approach in tobacco production as food source and smoke material – year 1976 and year 2000	81
<i>Une nouvelle approche de la production du tabac comme source alimentaire et matériau à fumer – an 1976 et an 2000</i>	
CHOUTEAU J., ALBO J.P.	
Etude in vitro des processus responsables des pertes de nicotine pendant la fermentation des tabacs humides	87
<i>In vitro study of the processes responsible for the loss of nicotine during fer- mentation of moist tobaccos</i>	
WAKEHAM H. R. R.	
Environmental carbon monoxide from cigarette smoking	93
<i>L'oxyde de carbone provenant de la fumée de cigarette, dans le milieu am- biant</i>	
ADAMS P. I.	
Changes in personal smoking habits brought about by changes in cigarette smoke yield	102
<i>Modifications des habitudes tabagiques des fumeurs en fonction de différents rendements en fumée des cigarettes</i>	

Abstracts – Résumés

SECTION A1

Mechanization and General

Mécanisation et Généralités

LABOUTIERE H., GABRIEL C.	
Développement de la mécanisation de la récolte du tabac en France de 1970 à 1976	113
<i>Development of mechanized tobacco harvesting in France from 1970 to 1976</i>	
NAITO T., KIMURA S.	
Mechanization for tobacco culture in Japan	114
<i>Mécanisation de la culture du tabac au Japon</i>	
SASAKI M., ASAI K., MIYAZONO T., SHIROZU A.	
Development of the transplanter for covered cultivation of tobacco in Japan	114
<i>Développement des machines à transplanter dans la culture du tabac sous couvert</i>	
BEUCHAT A., CAMMILLI A.	
La mécanisation de l'écimage, des traitements contre les bourgeons et de la récolte du tabac Bright: aspects techniques et économiques	115
<i>Mechanization of topping, bud treatment and Bright tobacco harvesting: technical and economic aspects</i>	
KNEISZL F., SCHIFFER L.	
Labor input in tobacco production with regard to 3 levels of mechaniza- tion	116

PRESIDENT OF CONGRESS

PRESIDENT DU CONGRES

Minomatsu Izumi

The Japan Tobacco & Salt Public Corporation

VICE-PRESIDENT

Kin-ichi Saito

The Japan Tobacco & Salt Public Corporation

President — Président

M. IZUMI

The Japan Tobacco & Salt Public Corporation

Members — Membres

S. ANDO

The Japan Tobacco & Salt Public Corporation

H. HARA

The Japan Tobacco & Salt Public Corporation

N. IMAGAWA

The Japan Tobacco & Salt Public Corporation

J. KATO

Mitsubishi Acetate Ltd.

S. MIYOKAWA

Fuji Flavor Co., Ltd.

K. NAKANISHI

Takasago Perfumery Co., Ltd.

T. SAKAYA

Daicel Ltd.

General Secretary — Secrétaire Général

E. TAMAKI

The Japan Tobacco & Salt Public Corporation

Chairman — Président

O. VON KIELMANSEGG

Generaldirektor-Stellvertreter der Austria Tabakwerke AG, Austria

Members — Membres

L. CREMONA

Direttore Generale, Amministrazione dei Monopoli di Stato, Italia

H. CULLMAN

President, Philip Morris International, New York, USA

N. HEDLUND

Vice-President, Swedish Tobacco Co., Sweden

M. ILTER

Directeur Général Adjoint, Turkish State Monopolies, Turkey

M. IZUMI

President, The Japan Tobacco & Salt Public Corporation

F. KRISTINUS

Vorsitzender des Verbandes der Cigarettenindustrie, Federal Republic of Germany

N. MANTZARIS

Directeur Général, Office National du Tabac, Grèce

P. MILLET

Directeur Général du S.E.I.T.A., France

Secretary — Secrétaire

P. RAY

Secrétaire Général

SCIENTIFIC COMMISSION

COMMISSION SCIENTIFIQUE

Chairman — Président

A. ARTHO F. J. Burrus & Cie., 2926 Boncourt, Switzerland

Vice-Chairmen — Vice-Présidents

L. McDONALD Tobacco Research Board of Rhodesia, P.O. Box 1909, Salisbury, Rhodesia

F. RESNIK Philip Morris Inc., 100 Park Avenue, New York, New York 10017, USA

Members — Membres

J. CHOUTEAU Institut Expérimental du Tabac, B.P. 168, 24108 Bergerac, France

R. CORBAZ Centre de Recherche sur le Tabac < SOTA >, Domaine de Changins, 1260 Nyon, Suisse

A. DEBAETS Faculté des Sciences Agronomiques, Coupure Links 533, B-9000 Gand, Belgique

C. DIETRICH Rinsoz et Ormond S.A., Rue du Collège 1, 1800 Vevey, Suisse

H. ELMENHORST Martin Brinkmann AG, Forschung und Entwicklung, 28 Bremen, Postfach 98, F.R.G.

F. HADDON Cigarette Components Ltd., 30 Abbey Estate, Mount Pleasant, Middx, Great Britain

G. HAYN Austria Tabakwerke AG., Postfach 14, A 1091 Wien, Austria

H. KUHN Austria Tabakwerke AG., Postfach 14, A 1091 Wien, Austria

A. LUBRANO Amministrazione dei Monopoli di Stato, Via della Luce 34 ab, Roma Italia

J. MERKER VEB Tabakkontor, 801 Dresden R.G. Weisseritzstrasse 3, G.D.R.

J. MICKOVSKI Institut Za Tutun, Prilep, Yougoslavia

P. SCHILTZ Institut Expérimental du Tabac, B.P. 168, 24108 Bergerac, France

J. SCHMIDT Landesanstalt für Tabakforschung, Forchheim/Bahnhof 7501, Kutscheweg 10, F.R.G.

M. SENKUS R. J. Reynolds Industries Inc., Winston-Salem, N.C. 27101, USA

K. SKULA Institut Expérimental de l'Industrie Tabacole, BAB pri Seredi, Tchécoslovaquie

A. SPEARS Lorillard Research Center, P.O. Box 21688, Greensboro, N.C. 27420, USA

D. VASSILIADIS Tobacco Institute of Greece, Drama, Greece

NOTICE
This document is
available from
the following source:

Environmental Carbon Monoxide from Cigarette Smoking

L'Oxyde de Carbone Provenant de la Fumée de Cigarette, dans le Milieu Ambiant

by Helmut R. R. WAKEHAM

Research Center, Philip Morris Inc., Richmond, Va., USA

SUMMARY

The current drive for bans on smoking in public places and in work places is based on the assertion that tobacco smoke in the atmosphere constitutes a health hazard to exposed non-smokers. It is said that the hazard arises mainly from carbon monoxide generated during the burning of a cigarette. Cigarette smoking is an insignificant source of carbon monoxide in the atmosphere as compared to other natural and man-made sources. Even in tightly closed spaces with a large percentage of smokers, only rarely is it possible to build-up concentrations which would exceed the established threshold limiting values for extended exposures. Examples including meeting rooms, houses, buses, and aircraft will be discussed. The resulting carboxyhemoglobin levels in nonsmokers will be considered. It is concluded that the carbon monoxide from environmental tobacco smoke is well below the amount needed to produce the maximum allowable limit of 4 % carboxyhemoglobin in the blood as suggested by the World Health Organization.

RÉSUMÉ

La campagne actuelle pour l'interdiction de fumer dans les lieux publics et de travail se base sur l'assertion que la fumée de tabac dans l'atmosphère constitue un danger pour la santé des non-fumeurs qui y sont exposés. On dit que le danger vient de l'oxyde de carbone engendré par la cigarette allumée.

La fumée de cigarette est une source insignifiante d'oxyde de carbone dans l'atmosphère comparée aux autres sources naturelles ou industrielles. Même dans des lieux clos, avec un pourcentage élevé de fumeurs, il n'est que rarement possible d'élever la concentration de CO à des niveaux dépassant ceux qui ont été établis comme valeurs-seuils pour l'exposition extensive. Parmi les exemples évoqués, on trouvera les salles de réunion, les maisons d'habitation, les autobus, et les avions. Les taux de carboxy-hémoglobine chez les non-fumeurs seront considérés. On estime, en conclusion, que l'oxyde de carbone dans le milieu, qui procède de la fumée de tabac, est bien en dessous de la quantité nécessaire pour produire les 4 %, considérés comme maximum par l'O.M.S., de carboxy-hémoglobine dans le sang.

INTRODUCTION

During the past decade the anticigarette movement has undergone a basic change in direction. Instead of attacking the smoker, it has now become the self-appointed protector of the

nonsmoker. In former times, even centuries ago, those opposed to smoking traditionally argued that smoking was harmful to the smoker. This viewpoint even had Judeo-Christian foundations and took on religious overtones. It was argued that in smoking (as well as

in drinking, taking drugs, eating unclean foods, or engaging in nonprocreative sex) your body. « the Temple of the living God », (1) was being defiled. Many elements of our modern free society have rejected this attitude. They have come to accept the notion that the individual is free to do as he pleases so long as he does not harm his neighbor.

Now, the anticigarette movement has cleverly seized upon this new approach to justify restricting the smoker. Cigarette smoke in the environment, the anticigarette activist says, is polluting the air so as to harm the nonsmoker. Smoking must be banned in all places where there is any possibility that smoke might be inhaled by innocent nonsmokers. Recently, there has even been proposed to the USA Federal Occupational Safety and Health Administration (2) that smoking should be banned in all *work places*. A *work place* is defined as a location where two or more persons are working. The basic rationale for thus restricting the freedom of the smoker is that in smoking he may be causing harm to someone else. The smoker is made to feel guilty in the presence of nonsmokers. He is even said to be mentally ill (or perhaps deranged) because of his insistence on hurting himself and others. The nonsmoker is being encouraged to bring pressure on the smoker to refrain from smoking.

This new move to protect the nonsmoker has brought forth a rash of studies to demonstrate how it might be that the nonsmoker is being harmed. By far the majority of these investigations is focused on carbon monoxide (CO) generated by the burning cigarette. In this paper I shall briefly examine those factors relating to the effects of environmental CO from cigarette smoking.

SOURCES OF CARBON MONOXIDE

It has been estimated that the earth's atmosphere at any given moment contains approximately 530 million metric tons of CO (3, 4) (Table 1). Natural and manmade sources produce annually ten times this amount. Because of the continual consumption of CO in a number of oxidation reactions in the atmosphere and stratosphere, and in microbial actions in the soil, the half life of atmospheric CO is only one-tenth of a year.

Note that cigarette smoking worldwide is estimated to produce only about 0.6 million tons of CO per year, an infinitesimal amount as compared to other sources. This estimate is based on an average figure of 0.100 gram per

TABLE 1

Annual production and losses
of atmospheric carbon monoxide (3, 4)

Production	Estimated production or loss in millions of tons
Atmospheric oxidation of methane	5,000.
Combustion of fossil fuels	270.
Escape from oceanic solution	150.
Tobacco smoking	0.6
Losses	
Atmospheric oxidation to CO ₂	4,500.
Stratospheric oxidation	500.
Soil microorganisms	500.
Equilibrium content	530.

cigarette for both mainstream and sidestream smokes. Clearly if we are to be concerned about CO from smoking, it must be from a local concentration point of view rather than from the total amount available in the atmosphere. For this reason, attention is usually focused on smoking in enclosed spaces such as rooms, buses, automobiles, airplanes, arenas, etc.

CONCENTRATIONS OF CO IN CLOSED SPACES

Carbon monoxide concentrations in nonurban outdoor atmospheres are usually less than 2 ppm. In highly populated cities average concentrations may range as high as 10 ppm (5), depending on proximity to sources (highways or streets, factories, etc.) and air circulation (or stagnation). In general, outdoor concentrations are less than the ambient air quality standard of 9 ppm or the USA Environmental Protection Agency standard limit of 8 ppm for 24-hour exposure (6).

In enclosed spaces such as homes, offices, elevators, workshops, autos, buses, aircraft, etc., maximum equilibrium concentrations of CO buildup from smoking depend on rates of cigarette smoking, volume of space, ventilation rates, and CO losses (7). A large number of combinations have been investigated and measured (8-11). Unfortunately, many of the tests were carried out under extreme conditions that

are completely unrealistic in terms of human exposures.

Some recent results are listed in Table 2. It is immediately evident from these data that under realistic conditions, CO concentrations from smoking are only one-fifth to one-tenth of those reported in sealed climatic chambers or unventilated rooms. Both RUSSELL (10) and HARKE (11, 12) found that in these closed, unventilated, smoke-filled rooms people experience annoying eye and nose irritations long before the CO concentration in the room reaches 40 ppm.

The American Conference of Governmental Industrial Hygienists (13) has established a Threshold Limiting Value for CO of 50 ppm as the maximum for daily 8-hour exposure. JONES and FAGAN (7) have shown that in a closed room or house it is almost impossible to attain this concentration of CO by smoking alone. The reason for this is that unless the space is tightly sealed, « normal » leakage through cracks around doors and windows permits substantial circulation of air into and out of the room (14), usually sufficient to change the total volume of air at least once every hour. There is, thus, an upper limit to the possible buildup of carbon monoxide concentration from smoking in the real-life situation. Many experimenters who

have studied the smoke-filled room have found this out, often to their surprise and dismay.

It is also of interest to note in this connection that the modern jet passenger aircraft is probably the most effectively ventilated enclosed space occupied by humans, since a complete change of air occurs every three minutes. Even if all the 165 passengers in a full Boeing 707 aircraft were smoking at the rate of four cigarettes per hour, the ambient carbon monoxide concentration would only increase by 12 ppm. The actual levels are much lower than this value (15).

In the arguments used to justify the imposition of laws restricting the rights of smokers to smoke only in restricted places, only those investigators who used extreme and unrealistic test conditions are most often cited. Exaggerated claims are made that CO from smoking greatly harms nearby nonsmokers. No one suggests that the best control for carbon monoxide from smoking in any indoor space is simply to provide reasonable ventilation, rather than restrict the freedom of the smoker (16).

INHALED CO AND CARBOXYHEMOGLOBIN

The main effect (if not the only effect) of


TABLE 2
Maximum CO concentrations recently reported in enclosed spaces

Situation	Estimated smoke generation rate Cigs/m ³ /hr.	Ventilation Air changes/hr.	Maximum CO ppm	Reference
<u>With little or no ventilation</u>				
Room 58 m ³	2.6	None (?)	50	(46)
Room 43 m ³	1.4	None (?)	38	(10)
Small autos on Hamburg streets	~3.0	Vents closed	24.3	(47)
Stationary auto	~3.0	Vents closed	110.4	(48)
Unventilated climate chamber 30 m ³	6.0	0.1	88 @ 10 min.	(49)
<u>With normal ventilation</u>				
Room 58 m ³	2.6	8.8	10	(46)
Party of 50 people in 170 m ³ room	0.26	7.0	7	(50)
Room 80 m ³	0.3	6.4	6	(51)
Small autos on Hamburg streets	~3.0	Vents open	12.0	(47)
Stationary auto in wind tunnel	~3	~60	5-6	(48)

inhaling CO is the formation of carboxyhemoglobin (COHb) in the blood (Figure 1). CO binds strongly to the hemoglobin preventing it from performing its intended function of transporting oxygen to the tissues and carbon dioxide to the lungs. The level of COHb (expressed as a percent of the hemoglobin so taken up) depends upon the concentration of CO in the lungs, the time of exposure, the rates of respiration and heart beat, exercise factors, etc. (17-21). Eventually under constant conditions an equilibrium concentration of COHb is established (22, 23). The formation of COHb is reversible so that at lower concentrations of CO, the COHb breaks down to release free hemoglobin again. Obviously, at high enough levels of COHb (50-70 %) the oxygen needed by various tissues is no longer supplied in adequate amounts and various adverse effects on body functions are observable. That such effects do not take place from inhaling cigarette smoke will be shown later.

Under these relationships between CO from smoke and COHb, the only important question is: How much COHb is formed in the exposed

Figure 1
Effects of high levels of carbon monoxide (56). Note that in the lower left hand corner are shown the values of 5 % COHb, the level found for the average smoker in an urban environment; and the atmospheric concentration value of 50 ppm, the Threshold Limiting Value established by the American Conference of Governmental Industrial Hygienists.



subjects? In fact this is the only complete statement one can make about CO exposure. Statements limited to transient concentrations of CO in smoke or in the atmosphere are inadequate to define what is taking place. Inferences from such marginal information about the effects of CO are usually based on unrealistic assumptions concerning the other conditions involved. For this reason, most of the information in the literature about CO concentrations from smoking is meaningless in terms of human experience.

OBSERVED COHb CONCENTRATIONS

It is impossible in the scope of this paper to review all the studies which have been made on the COHb concentrations in smokers and in nonsmokers. Such investigations are invariably complicated by the fact that differing CO concentrations are found in various environments even in the absence of smoking. In fact, the human body itself generates CO *de novo* so that the blood always contains from 0.4 to 1 % COHb (9, 24).

Table 3 summarizes some of the recent findings.

The pattern from these data is very clear. Nonsmokers will generally have less than 2 % COHb while smokers will average about 6 %, ranging from 2 to 13 % depending upon the extent and manner of smoking and on other environmental factors. Only in extreme situations such as those used by Russell in his experiments will nonsmokers experience COHb

TABLE 3
Carboxyhemoglobin Levels
in Smokers and Nonsmokers

Description	Mean (m) or Range (r) of % COHb	
	Nonsmokers	Smokers
UK pregnant women	m 1.1%	m 3.6%
Meat porters	m 1.6	m 5.1
Office workers (17)	m 1.3	m 6.2
London office workers (52)	m 1.12 r 0.1-2.7	m 5.5 r 2.2-13.0
29,000 USA blood donors (53)	m 1.39 r 0.4-6.9	m 5.57 r 0.8-11.9
3311 California longshoremen (54)	m 1.3	m 5.9
Munich population	m 2.36	m 7.38
Rural Bavarians (55)	m 1.03	m 6.06

levels in excess of 2 % from cigarette smoke-filled environments. Russell's (10) description of such an experiment is a classic :

"Twenty-one research and clerical colleagues volunteered to provide blood specimens before and after spending about 1 hour in a smoke-filled room. The conditions were deliberately made worse than would be likely to be encountered in natural social situations. The room was approximately 43 m³ (15 x 12 x 8 ft.). Ventilators were switched off and all windows were closed. Before the volunteers entered the room it was "smoked up" by leaving thirty cigarettes (" Piccadilly " tipped) to burn in ashtrays. During the experiment the smokers smoked thirty-two cigarettes and two cigars ; a further eighteen cigarettes were left to burn in ashtrays. During the average time of 78 minutes spent in the room the volunteers remained seated in the same place. The exposure was very unpleasant, causing eyes to burn and water. For most volunteers it was worse than they could recall having tolerated on normal social occasions."

Imagine, if you will, this experimental situation. The room had only 18 square meters of floor space, occupied by 21 persons. In 78

minutes smoke was generated from 80 cigarettes and two cigars. None was allowed to escape through ventilation. After all this, the maximum CO concentration in the room was only 38 ppm. For the nonsmokers the average COHb had increased from 1.6 % to 2.6 %. One wonders what the room conditions would have been even without smoking.

The World Health Organization (25) has suggested a maximum of 4 % COHb as a level which is safe from all possible adverse effects under continuous exposure conditions. In order to reach this level, the non-smokers in Russell's experiment would have had to remain in the intolerably smoky room at least another 78 minutes.

HARKE (11) carried out similar experiments and made the following concluding statement :

"It does not appear possible that such smoke densities can be generated by smokers. This experiment using smokers (rather than smoking machines) would only be reproducible when 50 persons were placed in the room (floor space 62 m²), and each of them would smoke three cigarettes during a period of 30 minutes. The irritation...would be so great that certainly a large majority of the smokers would have to leave the room prior to the end of the smoking




Figure 2
Temporal variations of COHb concentration (26). Note that the COHb level of the nonsmoking motor pool manager increases during the day when he is at work and quickly drops again at the end of the work day. The smoking housewife would have a daily average of about 5 % characteristic of smokers, the value shown in the lower left hand corner of Figure 1.

period."

It should be emphasized that because of the dynamic interchange of CO between the inhaled air and the COHb in the blood, the level of COHb is constantly changing, depending on circumstances. McILVAINE and associates (26) have measured COHb values at two-hour intervals throughout the day for smokers and nonsmokers. Some results are shown on Figure 2. Obviously, the definition of an average condition of exposure to carbon monoxide presents some difficulties.

HUMAN RESPONSE TO INCREASED COHb

Many investigators have carried out performance and pathological tests aimed at showing that increased COHb in the blood impairs the well being of the persons exposed to CO in the environment (27-29). Performance tests are generally directed to demonstrating the impairment of oxygen sensitive organs such as the eye, the brain, and the central nervous system. These researches indicate that no discernible effects occur in normal healthy adults below 10 % COHb. Some effects may be observed at relatively high and unusual COHb levels (15 % and over).

Recently, a number of studies have been carried out to explore the hypothesis that carbon monoxide exposure contributes to pathological changes leading to atherosclerosis(30). Pigeons(31), rabbits(32-34), dogs(34), primates(34-37), and man(38, 39) have all been investigated. No indications of any detrimental effects have so far been found at blood COHb levels of less than 15 %. The suggestion that carbon monoxide is implicated in the very weak statistical association between smoking and heart disease is very tenuous at best.

The recommendation by WHO of 4 % COHb as the maximum safe level must be considered as cautiously conservative. The evidence to date indicates that only if the individual is handicapped with severe heart disease might he have some difficulties at the 4 % COHb level (40, 41), but even then the probability is very low. Such a person may lack sufficient reserve to compensate for such loss in oxygen carrying capacity of the blood. Hence, if he were also exercising for a considerable period of time he might experience early onset of chest pains. (It is hardly likely that a room where people are smoking heavily is also a place where one would be engaging in strenuous exercise).

STEWART(42) in his overall review of the carbon monoxide and smoking situation has

published the information in Table 4 summarizing human responses to various levels of COHb (Table 4).

TABLE 4
Human Response to Elevated
Carboxyhemoglobin in the Blood (42)

% COHb	Healthy Adult	Heart Patients
0.3 - 0.7	Normal	Normal
1 - 5	Increased blood flow	May feel lack of oxygen
2 - 9	Exercise tolerance reduced	Chest pains with less exertion
16 - 20	Headache, lowered visual response	May be lethal for patients with severe cardiac problems
20 - 30	Throbbing headache, nausea, impaired manual dexterity	
30 - 40	Severe headache, nausea, and vomiting	
50 -	Coma, convulsions	
67 - 70	Lethal	

CONCLUSIONS

The following conclusion which appears in the Workshop Report entitled Environmental Tobacco Smoke Effects on the Non-Smoker published in the Scandinavian Journal of Respiratory Diseases (43), is supported by many investigators who have studied this problem(44, 45).

Concentrations that are present under realistic environmental conditions may reach about 10 ppm. It is true that higher concentrations have been reported, but they represent only transient values or levels reached under experimental conditions. If the exposure to 10 ppm were to prevail for 8 hours the resulting COHb concentration would be 1.9 %. This value is well below the suggested WHO maximum of 4 %. At these levels no adverse biological effects will occur in a population that can be expected to be present in such environments. It can thus be concluded that the carbon monoxide in environmental tobacco smoke does not represent a health hazard.

REFERENCES

1. The Bible, King James Translation. II Corinthian 6, 16.
2. ASH (Action on Smoking and Health) Petition to OSHA (Occupational Safety and Health Administration), October 8, 1974.
3. WEINSTOCK, B., and NIKI, H. (1972): Carbon monoxide balance in nature. *Science* 176: 290-292.
4. MAUGH, T. H., II. (1972): Carbon monoxide: Natural sources dwarf man's output. *Science* 177: 338-339.
5. EISENBUD, M., and EHRLICH, L. R. (1972): Carbon monoxide concentration trends in urban atmospheres. *Science* 176: 193-194.
6. Federal Register 36: 8187 (30 April 1971).
7. JONES, R. M., and FAGAN, R. (1974): Application of mathematical model for the buildup of carbon monoxide from cigarette smoking in rooms and houses. *ASHRAE Journal*: 49-53.
8. PENKALA, S. J., and de OLIVEIRA, G. (1975): The simultaneous analysis of carbon monoxide and suspended particulate matter produced by cigarette smoking. *Environ Res.* 9: 99-114.
9. GORBAN, C. M., KONDRAT'YEVA, I. I., and PODDUBNAYA, L. A. (1964): Gaseous activity products excreted by man when in an air tight chamber. In *Problems of Space Biology*. N. M. SISKAYAN and U. I. YAZDOVSKIY, Eds. Joint Publication Research Service, Washington, D. C.
10. RUSSELL, M. A., COLE, P. V., and BROWN, E. (1973): Absorption by non-smokers of carbon monoxide from room air polluted by tobacco smoke. *Lancet* 1: 576-579.
11. HARKE, H. P., BAARS, A., FRAHM, B., et al. (1972): Passive smoking. Concentration of smoke constituents in the air of large and small rooms as a function of number of cigarettes smoked and time. *Int. Arch. Arbeitsmed.* 29: 323-339.
12. HARKE, H. P. (1974): Air pollution in smoke-filled rooms. *Rev. Environ. Hlth.* 1: 305-326.
13. Threshold Limited Values for Airborn Contaminants. Adopted by ACGIH for 1972. In "American Conference of Governmental Industrial Hygienists." Cincinnati, Ohio.
14. HOEGG, U. R. (1972): Cigarette smoke in closed spaces. *Environ. Hlth. Perspect.* 2: 117-128.
15. Health Aspects of Smoking in Transport Aircraft (1971). Joint Study, Federal Aviation Administration, Department of Transportation and National Institute of Occupational Safety and Health, Department of Health, Education and Welfare.
16. McNALL, P. E., Jr. (1975): Practical methods of reducing airborne contaminants in interior spaces. *Arch. Environ. Hlth.* 30: 552-556.
17. HAWKINS, L. L. (1976): Blood carbon monoxide levels as a function of daily cigarette consumption and physical activity. *Brit. J. Indust. Med.* 33: 123-129.
18. LAWTHORP, P. J., and COMINS, B. T. (1970): Cigarette smoking and exposure to carbon monoxide. *Ann. N.Y. Acad. Sci.* 174: 135-147.
19. HOLLAND R. A. B. (1970): Reaction rates of carbon monoxide and hemoglobin. *Ann. N. Y. Acad. Sci.* 174: 154-171.
20. KLIMISCH, H. J., CHEVALIER, H. J., HARKE, H. P., et al. (1975): Uptake of carbon monoxide in blood of miniature pigs and other mammals. *Toxicology* 3: 301-310.
21. PETERSON, J. E. and STEWART, R. D. (1975): Predicting the carboxy-hemoglobin levels resulting from carbon monoxide exposures.
22. LANDAW, S. A. (1973): The effects of cigarette smoking on total body burden and excretion rates of carbon monoxide. *J. Occup. Med.* 15: 231-235.
23. DAVIES, D. M. (1975): The application of threshold limit values for carbon monoxide under conditions of continuous exposure. *Ann. Occup. Hygiene* 18: 21-28.
24. Processes that influence the body carbon monoxide stores. Part I., in *Biological Effects of Carbon Monoxide*. (1970) R. F. COBURN, Ed. *Ann. N. Y. Acad. Sci.* 174: 5-75.
25. Air Quality Criteria and Guides for Urban Air Pollutants (1972). WHO (World Health Organization) technical Rep. Ser. #506.
26. McILVAINE, P. M., NELSON, W. C., and BARTLETT, D., Jr. (1969): Temporal variation of carboxyhemoglobin concentrations. *Arch. Environ. Hlth* 19: 83-91.
27. For details of some of these the reader is referred to Parts II and IV of *Biological Effects of Carbon Monoxide*, *Ann. N. Y. Acad. Sci.* 174: 148-429, 1970, and *Environmental Tobacco Smoke Effects on the Non-Smoker: Report from a Workshop*, R. Rylander, Ed., *Scand. J. Resp. Dis., Suppl.* 91, 1974.
28. McFARLAND, R. A., FORBES, W. H., STROUD, H. W., et al. (1973): A study of

- the effects of low levels of carbon monoxide upon humans performing driving tasks. Harvard School of Public Health, Boston, Mass.: 88.
29. WEBER, A., JERMINI, C., and GRANDJEAN, E. (1975): Effects of low carbon monoxide concentrations on flicker fusion frequency and on subjective feelings. *Int. Arch. Occup. Environ. Hlth.* 36: 87-103.
 30. WALD, N., HOWARD, S., SMITH, P. G., et al. (1973): Association between atherosclerotic diseases and carboxyhaemoglobin levels in tobacco smokers. *Brit. Med. J.* 1: 761-765.
 31. ARMITAGE, A. K., DAVIES, R. F., and TURNER, D. M. (1976): The effects of carbon monoxide on the development of atherosclerosis in the White Carneau pigeon. *Atherosclerosis* 23: 333-344.
 32. THOMSEN, H. and KJELDSEN, K. (1975): Aortic intimal injury in rabbits. An evaluation of a threshold limit. *Arch. Environ. Hlth.* 30: 604-607.
 33. THOMSEN, H. and KJELDSEN, K. (1975): Threshold limit for carbon monoxide-induced myocardial damage. An electron microscopic study in rabbits. *Arch. Environ. Hlth.* 29: 73-78.
 34. ASTRUP, P. (1974): Tobacco smoking and coronary disease. *Acta Cardiol. Suppl.* 20: 105-117.
 35. MALINOW, M. R., McLAUGHLIN, P., DHINDSA, D. S., et al. (1976): Failure of carbon monoxide to induce myocardial infarction in cholesterol-fed Cynomolgus monkeys. *Cardiovascular Res.* 10: 101-108.
 36. THOMSEN, H. (1974): Carbon monoxide-induced atherosclerosis in primates. *Atherosclerosis* 20: 233-240.
 37. ECKARDT, R. E., MACFARLAND, H. N., ALARIE, Y. C. E., et al. (1972): The biologic effect from long-term exposure of primates to carbon monoxide. *Arch. Environ. Hlth.* 25: 381-387.
 38. MASTER, K. (1976): Carbon monoxide, atherosclerosis and natives of the Highlands of New Guinea. *New Eng. J. Med.* 294: 557.
 39. PARVING, H. H. (1972): The effect of hypoxia and carbon monoxide exposure on plasma volume and capillary permeability to albumin. *Scand. J. Clin. Lab. Invest.* 30: 49-56.
 40. ARONOW, W. S. and ROKAW, S. N. (1971): Carboxyhemoglobin caused by smoking nonnicotine cigarettes. Effects in angina pectoris. *Circulation* 44: 782-788.
 41. ARONOW, W. S., CASSIDY, J., VANGROW, J. S., et al. (1974): Effect of cigarette smoking and breathing carbon monoxide on cardiovascular hemodynamics in arterial patients. *Circulation* 50: 340-347.
 42. STEWART, R. D. (1974): The effects of low concentrations of carbon monoxide in man, in *Environmental Tobacco Smoke Effects on the Non-Smoker. Report from a Workshop*. R. Rylander, Ed. *Scand. J. Resp. Dis., Suppl.* 91: 56-62.
 43. RYLANDER, R. (1974): Perspectives on environmental tobacco smoke effects, in *Environmental Tobacco Smoke Effects on the Non-Smoker. Report from a Workshop*. R. Rylander, Ed. *Scand. J. Resp. Dis., Suppl.* 91: 79-87.
 44. ANTWEILER, H. (1975): Passive smoking causes no health risk by CO, because the COHb-value, even in extreme situations, does not exceed 2.6%. *Arbeitsmed. Sozialmed. Präventivmed.* 10: 245-248.
 45. ARONOW, W. S. Effect of passive smoking on the cardiovascular and respiratory systems. Presented at the 3rd World Conference on Smoking & Health, Waldorf Astoria Hotel, New York, NY, June 2-5, 1975.
 46. HARKE, H. P. (1970): The problem of Passive Smoking. *Munch. Med. Wochenschr* 112: 1328-2334.
 47. HARKE, H. P. and PETERS, H. (1974): The problem of passive smoking. III. The influence of smoking on the CO concentration in driving automobiles. *Int. Arch. Arbeitsmed.* 33: 221-229.
 48. HARKE, H. P., LIEDL, W., and DENKER, D. (1974): The problem of passive smoking. II. Investigations of CO level in the automobile after cigarette smoking. *Int. Arch. Arbeitsmed.* 33: 207-220.
 49. JERMINI, C., WEBER, A., and GRANDJEAN, E. (1976): Quantitative determination of various gas-phase components of the side-stream smoke of cigarettes in the room air as a contribution to the problem of passive-smoking. *Int. Arch. Occup. Environ. Hlth.* 36: 169-181.
 50. BRIDGE, D. P. and CORN, M. (1972): Contribution to the assessment of exposure of non-smokers to air pollution from cigarette and cigar smoke in occupied spaces. *Environ. Res.* 5: 192-209.
 51. ANDERSON, G. and DALHAMM, T. (1973): Health risks due to passive smoking. *Läkartidn.* 70: 2833-2836.
 52. COLE, P. V. (1975): Comparative effects of atmospheric pollution and cigarette smoking on carboxyhaemoglobin levels in man.

- Nature 255: 699-701.
53. STEWART, R.D., BARETTA, E.D., PLATTE, L.R., et al. (1974): Carboxyhemoglobin levels in American blood donors. JAMA 229: 1187-1195.
54. GOLDSMITH, J.R. and LANDAW, S.A. (1968): Carbon monoxide and human health, Science 162: 1352-1359.
55. SCHEIDEMANDEL, V. and DAUM, S. (1973): Carboxyhemoglobin concentrations of the Munich population. Münch. Med. Wochenschr. 115: 109-112.
56. FORBES, W.H. (1970): Carbon monoxide uptake via the lungs. Ann. N.Y. Acad. Sci. 174: 72-75.

The author's address:
Philip Morris Inc., Research Center, P.O. Box
26583, Richmond, Va., USA.

DEBATE

KRUSZYNSKI. Dr. Harke has published a study on changes in skin temperatures for active and passive smokers in extreme room conditions. I have not seen any confirmation of these findings anywhere. Have you seen a confirmation of

Harke's findings or made own experiments on that item?

WAKEHAM. I know of no investigation to confirm Harke's findings.